

GENUS 2 FIELDS WITH DEGREE 3 ELLIPTIC SUBFIELDS

T. SHASKA

ABSTRACT. In this paper we study genus 2 function fields K with degree 3 elliptic subfields. We show that the number of $\text{Aut}(K)$ -classes of such subfields of fixed K is 0,1,2 or 4. Also we compute an equation for the locus of such K in the moduli space of genus 2 curves.

1. INTRODUCTION

We study genus two curves $\mathcal{C}$ whose function fields have a degree 3 subfield of genus 1. Such subfields we call elliptic subfields. Such curves $\mathcal{C}$ have already occurred in the work of Hermite, Goursat, Burkhardt, Brioschi, and Bolza, see Krazer [6] (p. 479). More generally, degree n elliptic subfields of genus 2 fields have been studied by Frey [1], Frey and Kani [2], Kuhn [8], Shaska [11], Shaska and Voelklein [13]. In the degree 3 case, explicit equations can be used to answer questions which remain open in the general case. This was the theme of the author's PhD thesis (see [12]) which also covered the case of degree 2 (which is subsumed in [13]).

Equation (5) gives a normal form for pairs (K, E) where E is a degree 3 elliptic subfield of K . This normal form depends on two parameters $a, b \in k$. Isomorphism classes of pairs (K, E) are parameterized by parameters u, v where $u := ab$ and $v := b^3$. This yields an expression for the invariants (i_1, i_2, i_3) of K (see Igusa [9]) in terms of u and v . Our central object of study is the map

$$\theta : (u, v) \rightarrow (i_1, i_2, i_3)$$

This map is shown to have degree 2, which means that a general genus 2 field having elliptic subfields of degree 3 has exactly two such subfields. There are exactly four genus 2 fields which have four $\text{Aut}(K)$ -classes and a 1-dimensional family of such fields with one $\text{Aut}(K)$ -class of such subfields.

Although the map θ is given by explicit equations (see (13)), it is a non-trivial computational task to show that it has degree 2 and to compute the action of its Galois group on u and v . The direct approach exceeds available computer power, so we introduce auxiliary parameters r_1, r_2 that parameterize pairs of cubic polynomials. They arise from the fact that the subfield E induces a particular sextic defining K which splits naturally as a product of 2 cubic polynomials. We show that θ factorizes as

$$(u, v) \rightarrow (r_1, r_2) \rightarrow (i_1, i_2, i_3)$$

where the latter map is birational, and the former has degree 2. Thus r_1, r_2 yield a birational parameterization of the locus $\mathcal{L}_3$ of genus 2 fields having a degree 3 elliptic subfield. That $\mathcal{L}_3$ is a rational variety follows also from the general theory of "diagonal modular surfaces", see Kani [4]. We also compute the equation in i_1, i_2, i_3 that defines $\mathcal{L}_3$ as a sublocus of the moduli space $\mathcal{M}_2$ of genus 2 curves.

We further find relations between the j -invariants of the degree 3 elliptic subfields of K and classify all K in $\mathcal{L}_3$ with extra automorphisms.

All the computations were done using Maple [10].

Acknowledgment: I would like to express my sincere gratitude to my PhD advisor Prof. H. Voelklein for all the time and effort spent in guiding me towards my dissertation (from which this paper originated).

2. GENUS TWO FIELDS WITH DEGREE 3 ELLIPTIC SUBFIELDS

Let k be an algebraically closed field of characteristic 0. All function fields will be over k . Let K be a genus 2 function field and E a degree 3 elliptic subfield of K .

The associated extension $k(X)/k(U)$. It is well-known that K has exactly one genus zero subfield of degree 2, which we denote by $k(X)$. The generator of $\text{Gal}(K/k(X))$ is the hyperelliptic involution of K . It fixes each elliptic subfield of K , see Tamme [14]. Hence the field $E \cap k(X)$, denote it by $k(U)$, is a subfield of E of degree 2.

$$\begin{array}{ccc} K & \xrightarrow{2} & k(X) \\ 3 \mid & & \mid 3 \\ E & \xrightarrow{2} & k(U) \end{array}$$

Ramification of K/E . Either K/E is ramified at exactly 2 places of K , of ramification index 2, or at one place of K , of ramification index 3. It follows immediately from the Riemann-Hurwitz formula. The former (resp., latter) case we call the non-degenerate (resp., degenerate) case, as in [1].

Definition 1. A **non-degenerate pair** (resp., **degenerate pair**) is a pair (K, E) such that K is a genus 2 field with a degree 3 elliptic subfield E where the extension K/E is ramified in two (resp., one) places. Two such pairs (K, E) and (K', E') are called isomorphic if there is a k -isomorphism $K \rightarrow K'$ mapping $E \rightarrow E'$.

Ramification of $k(X)/k(U)$. In the non-degenerate (resp., degenerate) case $k(X)/k(U)$ is ramified at exactly 4 (resp., 3) places of $k(X)$ each of ramification index 2 (resp., one of index 3 and the other two of index 2), see [1].

Invariants of K . We denote by J_2, J_4, J_6, J_{10} the classical invariants of K , for their definitions see [9] or [13]. These are homogeneous polynomials (of degree indicated by subscript) in the coefficients of a sextic $f(X, Z)$ defining K

$$Y^2 = f(X, Z) = a_6 X^6 + a_5 X^5 Z + \cdots + a_1 X Z^5 + a_0$$

and they are a complete set of $SL_2(k)$ -invariants (acting by coordinate change). The absolute invariants

$$(1) \quad i_1 := 144 \frac{J_4}{J_2^2}, \quad i_2 := -1728 \frac{J_2 J_4 - 3 J_6}{J_2^3}, \quad i_3 := 486 \frac{J_{10}}{J_2^5}$$

are even $GL_2(k)$ -invariants. Two genus 2 curves with $J_2 \neq 0$ are isomorphic if and only if they have the same absolute invariants.

Main Theorem: Let K be a genus 2 field and $e_3(K)$ the number of $\text{Aut}(K/k)$ -classes of elliptic subfields of K of degree 3. Then;

- i) $e_3(K) = 0, 1, 2$, or 4
 ii) $e_3(K) \geq 1$ if and only if the classical invariants of K satisfy the irreducible equation $F(J_2, J_4, J_6, J_{10}) = 0$ displayed in Appendix A.

There are exactly two genus 2 curves (up to isomorphism) with $e_3(K) = 4$, see 4.2. The case $e_3(K) = 1$ (resp., 2) occurs for a 1-dimensional (resp., 2-dimensional) family of genus 2 curves, see section 4.

2.1. The non-degenerate case. Let (K, E) be a non-degenerate pair and $k(X)$ and $k(U)$ their associated genus 0 subfields. Both $k(X)/k(U)$ and $E/k(U)$ are ramified at 4 places of $k(U)$, three of which are in common. Take the common places to be $U = q_1, q_2, q_3$. Also, take $U = 0$ (resp., $U = \infty$) the place ramified in $k(X)$ but not in E (resp., in E but not $k(X)$). Take $X = 0$ (resp., $X = \infty$) the place over $U = 0$ of ramification index 2 (resp., 1). In the following figure bullets (resp., circles) represent places of ramification index 2 (resp., 1).




FIGURE 1. Ramification of $k(X)/k(U)$

Thus, $U = \phi(X) = l \frac{X^2}{X^3 + aX^2 + bX + c}$, where $l, a, b, c \in k$ such that $X^3 + aX^2 + bX + c$ has no multiple roots and $l, c \neq 0$. We can make $l = 1$ and $c = 1$ by replacing U and X by scalar multiples. Then,

$$(2) \quad U = \phi(X) = \frac{X^2}{X^3 + aX^2 + bX + 1}$$

Remark 2. These conditions determine X up to multiplication by a third root of unity $\xi_3 \in k$. Replacing X by $\xi_3 X$ replaces (a, b) by $(\xi_3^2 a, \xi_3 b)$. Invariants of this transformation are:

$$(3) \quad \begin{aligned} u &= ab \\ v &= b^3 \end{aligned}$$

The derivative of $\phi(X)$ is

$$\phi'(X) = -\frac{X(X^3 - bX - 2)}{(X^3 + aX^2 + bX + 1)^2}$$

Taking the resultant with respect to z of $z^2 - bz - 2$ and $\frac{\phi(x) - \phi(z)}{x - z}$ we get

$$(4) \quad (x^3 - bx - 2)(4x^3 + b^2x^2 + 2bx + 1)$$

The roots of this polynomial correspond to the 6 places of $k(X)$ over the places $U = q_1, q_2, q_3$. Thus, the 6 places of $k(X)$ ramified in K correspond to the roots of the polynomial on the right side of

$$(5) \quad Y^2 = (X^3 + aX^2 + bX + 1)(4X^3 + b^2X^2 + 2bX + 1)$$

which gives an equation of K . The discriminant of the sextic is nonzero, hence

$$(6) \quad \Delta := (4a^3 + 27 - 18ab - a^2b^2 + 4b^3)^2(b^3 - 27) \neq 0$$

One checks that the element

$$(7) \quad V = Y \frac{X^3 - bX - 2}{F(X)^2}$$

satisfies

$$V^2 = (U - q_1)(U - q_2)(U - q_3) = U^3 + 2 \frac{ab^2 - 6a^2 + 9b}{R} U^2 + \frac{12a - b^2}{R} U - \frac{4}{R}$$

where $R := \frac{\Delta}{(b^3 - 27)}$. Thus, $E = k(U, V)$.

Lemma 3. *For $(a, b) \in k^2$ with $\Delta \neq 0$, equation (5) defines a genus 2 field $K_{a,b} = k(X, Y)$. It has a non-degenerate degree 3 elliptic subfield $E_{a,b} = k(U, V)$, where U and V are given in (2) and (7). Two such pairs $(K_{a,b}, E_{a,b})$ and $(K_{a',b'}, E_{a',b'})$ are isomorphic if and only if $u = u'$ and $v = v'$ (where u, v and u', v' are associated with a, b and a', b' , respectively, by (3)).*

Proof. The first two statements follow by reversing the above arguments. If $u = u'$, $v = v'$ then $a' = \xi_3^i a$, $b' = \xi_3^i b$ for some i . Then clearly the two associated non-degenerate pairs are isomorphic. The converse follows from Remark 2. $\square$

Proposition 4. *The $(u, v) \in k^2$ with $\Delta \neq 0$ bijectively parameterize the isomorphism classes of non-degenerate pairs (K, E) (via the parameterization defined in Lemma 3).*

Proof. The proof follows from Lemma 3. $\square$

From the normal form of K in (5) we can compute the classical invariants J_2, J_4, J_6, J_{10} in terms of u, v . These expressions satisfy equation (33) which proves one implication of claim ii) of the theorem for the non-degenerate pairs. In section 4 we explain how equation (33) was found.


2.2. The degenerate case. Let K/E be a degenerate pair. Then $k(X)/k(U)$ is ramified at exactly three places of $k(U)$ which are also ramified in E . Let $U = 0$ be the place ramified in E but not in $k(X)$. Take $U = \infty$ be the place of $k(U)$ that is totally ramified in $k(X)$ and $X = \infty$ the place over it.

Then, $U = l(X^3 + aX^2 + bX + c)$ for $a, b, c, l \in k$ such that $X^3 + aX^2 + bX + c$ has no multiple roots and $l \neq 0$. Replacing X by $X + \frac{a}{3}$ and U by a scalar multiple we get $U = X^3 + bX + c$. Note that $b \neq 0$ (otherwise two of the places of $k(U)$ ramified in $k(X)$ coalesce). Thus, by replacing X by a scalar multiple we can also get $b = 1$. Then,

$$(8) \quad U = X^3 + X + c$$

As in Lemma 3 we get $K = k(X, Y)$ such that

$$(9) \quad Y^2 = (3X^2 + 4)(X^3 + X + c)$$

FIGURE 2. Ramification of $k(X)/k(U)$, degenerate case.

where $c^2 \neq -\frac{4}{27}$. Then, $V = Y(3X^2 + 1)$ satisfies the equation

$$V^2 = U(27U^2 - 54cU + 4 + 27c^2)$$

Thus, $E = k(U, V)$.

Remark 5. In this case, those $w := c^2$ with $w \neq -\frac{4}{27}$ bijectively parameterize the isomorphism classes of degenerate pairs K/E . The proof is analogous to the non-degenerate case.

Since

$$\begin{aligned} J_2 &= 774 \\ J_4 &= 36(268 + 837w) \\ J_6 &= 36(76760 + 290574w - 729w^2) \\ J_{10} &= 432(27w + 4) \end{aligned} \tag{10}$$

the map

$$\begin{aligned} k \setminus \left\{ -\frac{4}{27} \right\} &\rightarrow k^3 \\ w &\rightarrow (i_1, i_2, i_3) \end{aligned}$$

is injective. Thus, a genus 2 field K has at most one $\text{Aut}(K)$ -orbit of subfields E such that (K, E) is a degenerate pair. One can check that J_2, J_4, J_6, J_{10} satisfy equation (33). This completes the proof that the condition in part (ii) of Main Theorem is necessary.

From (10) we compute that the locus of those K being part of a degenerate pair (K, E) is given by

$$\begin{aligned} 341880i_1^2 - 2550732480i_1 + 4023336960 + 611249816i_2 &= 0 \\ 161446939560824832i_3 + 23630752512i_1^2 - 6321363049i_1^3 - 29445783552i_1 \\ + 12230590464 &= 0 \end{aligned} \tag{11}$$

(Note $J_2 = 774 \neq 0$ on this locus, so i_1, i_2, i_3 are everywhere defined). For later use we note the following: If (K, E_0) is a non-degenerate pair with parameters u, v then there exists a degenerate pair (K, E_1) (for same K) if and only if

$$2v - 9u + 27 = 0 \tag{12}$$

This is obtained by expressing i_1, i_2, i_3 in (11) in terms of u, v .

3. FUNCTION FIELD OF $\mathcal{L}_3$

The absolute invariants i_1, i_2, i_3 in terms of u, v are

(13)

$$\begin{aligned} i_1 &= \frac{144}{v(-405 + 252u + 4u^2 - 54v - 12uv + 3v^2)^2} (1188u^3 - 8424uv + u^4v - 24u^4 \\ &\quad + 14580v - 66u^3v + 138uv^2 + 297u^2v + 945v^2 - 36v^3 + 9u^2v^2) \\ i_2 &= -\frac{864}{v^2(-405 + 252u + 4u^2 - 54v - 12uv + 3v^2)^3} (-81v^3u^4 + 2u^6v^2 + 234u^5v^2 \\ &\quad + 3162402uv^2 - 21384v^3u + 26676v^4 - 473121v^3 - 72u^6v - 5832v^4u + 14850v^3u^2 \\ &\quad - 72v^3u^3 + 324v^4u^2 - 650268u^3v - 5940u^3v^2 - 3346110v^2 + 432u^6 - 1350u^4v^2 \\ &\quad + 136080u^4v - 7020u^5v - 307638u^2v^2) \\ i_3 &= -243 \frac{(v-27)(4u^3 - u^2v - 18uv + 4v^2 + 27v)^3}{v^3(-405 + 252u + 4u^2 - 54v - 12uv + 3v^2)^5} \end{aligned}$$

We will sometimes view u, v as parameters from k and sometimes as the corresponding coordinate functions on k^2 . From the context it will be clear which point of view we are taking.

Lemma 6. $[k(i_1, i_2, i_3) : k(u, v)] \leq 2$

Proof. From the resultants of equations in (13) we determine that $[k(v) : k(i_1, i_2)] = 16$, $[k(v) : k(i_2, i_3)] = 40$, and $[k(v) : k(i_1, i_3)] = 26$. We can show that $u \in k(i_1, i_2, i_3, v)$, the expression is large and we don't display it (the interested reader can check [12]). Since $[k(u, v) : k(i_1, i_2, i_3)]$ must be a common divisor of 16, 26, and 40, then the claim follows. $\square$

We need to show that $[k(u, v) : k(i_1, i_2, i_3)] = 2$. Since we know that $[k(u, v) : k(i_1, i_3)] = 26$, this is equivalent to showing that $[k(i_1, i_2, i_3) : k(i_1, i_3)] = 13$. This follows once we have equation (33) in terms of i_1, i_2, i_3 but at this stage we cannot derive it: Eliminating u, v from equations (13) exceeds available computer power. We use additional invariants r_1, r_2 to overcome this problem.

3.1. Invariants of Two Cubics. We define the following invariants of two cubic polynomials. For $F(X) = a_3X^3 + a_2X^2 + a_1X + a_0$ and $G(X) = b_3X^3 + b_2X^2 + b_1X + b_0$ define

$$H(F, G) := a_3b_0 - \frac{1}{3}a_2b_1 + \frac{1}{3}a_1b_2 - a_0b_3$$

We denote by $R(F, G)$ the resultant of F and G and by $D(F)$ the discriminant of F . Also,

$$r_1(F, G) = \frac{H(F, G)^3}{R(F, G)}, \quad r_2(F, G) = \frac{H(F, G)^4}{D(F)D(G)}$$

Remark 7. In [7] it is shown that r_1, r_2 , and $r_3 = \frac{H(F, G)^2}{J_2(F, G)}$ form a complete system of invariants for unordered pairs of cubics.

For $F(X) = X^3 + aX^2 + bX + 1$ and $G(X) = 4X^3 + b^2X^2 + 2bX + 1$ and u, v as in (3) we have

$$(14) \quad \begin{aligned} r_1(F, G) &= 27 \frac{v(v-9-2u)^3}{4v^2 - 18uv + 27v - u^2v + 4u^3} \\ r_2(F, G) &= -1296 \frac{v(v-9-2u)^4}{(v-27)(4v^2 - 18uv + 27v - u^2v + 4u^3)} \end{aligned}$$

Taking the resultants from the above equations we get the following equations for u and v over $k(r_1, r_2)$:

$$(15) \quad \begin{aligned} &65536r_1r_2^3u^2 + (42467328r_2^4 + 21233664r_2^4r_1 + 480r_2r_1^4 + 2r_1^5 + 41472r_2^2r_1^3 \\ &+ 1548288r_2^3r_1^2 - 294912r_2^3r_1)u - 382205952r_2^4 + 238878720r_2^4r_1 - 2654208r_2^3r_1 \\ &+ 13934592r_2^3r_1^2 + 285696r_2^2r_1^3 + 2400r_2r_1^4 + 7r_1^5 = 0 \end{aligned}$$

$$(16) \quad \begin{aligned} &16384v^2r_2^3 + (221184r_2^3r_1 + r_1^4 + 11520r_2^2r_1^2 - 442368r_2^3 + 192r_2r_1^3)v \\ &- 5971968r_2^3r_1 - 864r_2r_1^3 - 124416r_2^2r_1^2 - 2r_1^4 = 0 \end{aligned}$$

Roots of equation (15) (resp., equation (16)) are u and $\beta(u)$ (resp., v and $\beta(v)$) where,

$$(17) \quad \begin{aligned} \beta(u) &= \frac{(v-3u)(324u^2 + 15u^2v - 378uv - 4uv^2 + 243v + 72v^2)}{(v-27)(4u^3 + 27v - 18uv - u^2v + 4v^2)} \\ \beta(v) &= -\frac{4(v-3u)^3}{4u^3 + 27v - 18uv - u^2v + 4v^2} \end{aligned}$$

It follows that either $[k(u, v) : k(r_1, r_2)] = 2$ or $k(u, v)/k(r_1, r_2)$ is Galois with Klein 4-group as Galois group. In the latter case, the map

$$(18) \quad \begin{aligned} \tau : k(u, v) &\rightarrow k(u, v) \\ u &\rightarrow u \\ v &\rightarrow \beta(v) \end{aligned}$$

is an involutory automorphism of $k(u, v)$. But by plugging in equation (17) we see that it is not involutory. Thus, $[k(u, v) : k(r_1, r_2)] = 2$ and $Gal_{k(u, v)/k(r_1, r_2)} = \langle \beta \rangle$.

Lemma 8. *The fields $k(i_1, i_2, i_3)$ and $k(r_1, r_2)$ coincide, hence $[k(u, v) : k(i_1, i_2, i_3)] = 2$. Moreover;*

$$(19) \quad \begin{aligned} i_1 &= \frac{9}{4} \frac{(13824r_1^3r_2^2 + 442368r_1^2r_2^3 + 5308416r_1r_2^4 + 192r_1^4r_2 + r_1^5 + 786432r_1r_2^3 + 9437184r_2^4)}{r_1(-1152r_2^2 + 96r_2r_1 + r_1^2)^2} \\ i_2 &= \frac{27}{8r_1^2(-1152r_2^2 + 96r_2r_1 + r_1^2)^3} (+79626240r_1^4r_2^4 - 4076863488r_1^2r_2^5 + 34560r_1^6r_2^2 \\ &+ 12230590464r_1^2r_2^6 + 32614907904r_1r_2^6 + 14495514624r_2^6 + 288r_1^7r_2 + 2211840r_1^5r_2^3 \\ &+ r_1^8 - 212336640r_1^3r_2^4 + 1528823808r_1^3r_2^5 - 2359296r_1^4r_2^3) \\ i_3 &= -521838526464 \frac{r_2^9}{r_1^2(-1152r_2^2 + 96r_2r_1 + r_1^2)^5} \end{aligned}$$

Proof. Equations (19) are verified by expressing all variables in terms of u, v (These equations were found by taking suitable resultants). It follows that $k(i_1, i_2, i_3) \subset$

$k(r_1, r_2)$. Equality follows since $[k(u, v) : k(r_1, r_2)] = 2$ and $[k(u, v) : k(i_1, i_2, i_3)] \leq 2$ (see Lemma 6). $\square$

Remark 9. To find equation (33) we eliminate r_1 and r_2 from equations (19). The resulting equation (33) has degree 8, 13, and 20 in i_1, i_2, i_3 respectively.

4. PROOF OF MAIN THEOREM

The map

$$\theta : (u, v) \rightarrow (i_1, i_2, i_3)$$

given by (13) has degree 2, by previous section and it is defined when $J_2 \neq 0$. For now we assume that $J_2 \neq 0$ (The case $J_2 = 0$ is treated in section 4.2). Denote the minors of the Jacobian matrix of θ by $M_1(u, v), M_2(u, v), M_3(u, v)$. The solutions of

$$(20) \quad \begin{cases} M_1(u, v) = 0 \\ M_2(u, v) = 0 \\ M_3(u, v) = 0 \end{cases}$$

consist of the (non-singular) curve

$$(21) \quad 8v^3 + 27v^2 - 54uv^2 - u^2v^2 + 108u^2v + 4u^3v - 108u^3 = 0$$

and 7 isolated solutions which we display in table (1) together with the corresponding values (i_1, i_2, i_3) and properties of the corresponding genus 2 field K .

(u, v)	(i_1, i_2, i_3)	$Aut(K)$	$e_3(K)$
$(-\frac{7}{2}, 2)$	$J_{10} = 0$, no associated genus 2 field K		
$(-\frac{775}{8}, \frac{125}{96}),$ $(\frac{25}{2}, \frac{250}{9})$	$-\frac{8019}{20}, -\frac{1240029}{200}, \frac{531441}{100000}$	D_4	2
$(27 - \frac{77}{2}\sqrt{-1}, 23 + \frac{77}{9}\sqrt{-1}),$ $(27 + \frac{77}{2}\sqrt{-1}, 23 - \frac{77}{9}\sqrt{-1})$	$(\frac{729}{2116}, \frac{1240029}{97336}, \frac{531441}{13181630464})$	D_4	2
$(-15 + \frac{35}{8}\sqrt{5}, \frac{25}{2} + \frac{35}{6}\sqrt{5}),$ $(-15 - \frac{35}{8}\sqrt{5}, \frac{25}{2} - \frac{35}{6}\sqrt{5})$	$81, -\frac{5103}{25}, -\frac{729}{12500}$	D_6	2

TABLE 1. Exceptional points where $\det(Jac(\theta)) = 0$

Let $\bar{u}, \bar{v}, \bar{i}_1, \bar{i}_2, \bar{i}_3$ denote the restrictions of the corresponding functions to the curve (21). Eliminating u from (21) and the defining equation for i_1 in (13) we get a relation between v and i_1 which shows $[k(\bar{v}, \bar{i}_1) : k(\bar{i}_1)] = 8$; also we get $\bar{u}$ expressed as a rational function of $\bar{v}$ and $\bar{i}_1$. Hence $k(\bar{v}, \bar{i}_1) = k(\bar{u}, \bar{v})$ and so $[k(\bar{u}, \bar{v}) : k(\bar{i}_1)] = 8$. Similarly we get a relation between $\bar{v}$ and $\bar{i}_2$. Eliminating $\bar{v}$ gives

$$(22) \quad G(\bar{i}_1, \bar{i}_2) = 0$$

an irreducible equation between $\bar{i}_1$ and $\bar{i}_2$ of degree 8 (resp., 12) in $\bar{i}_1$ (resp., $\bar{i}_2$). It is fully displayed as equation (34) in Appendix B. Thus, $[k(\bar{i}_1, \bar{i}_2) : k(\bar{i}_1)] = 8$. Since also $[k(\bar{u}, \bar{v}) : k(\bar{i}_1)] = 8$, it follows that $k(\bar{u}, \bar{v}) = k(\bar{i}_1, \bar{i}_2)$. Hence, $e_3(K) = 1$ for any K such that the associated u and v satisfy equation (21) except possibly those (u, v) that lie over singular points of the curve (22). We check that all these singular points have multiplicity 2, hence there are at most two (u, v) points over each of them.

Remark 10. *If K has one subfield of degree 3, then it has at least two distinct such subfields. This follows from the explicit construction in [12], Lemma 5.4. (It is actually true for degree n elliptic subfields, for any n , see [13]). Since $e_3(K) = 1$ for a generic curve K of the locus (22), such K has $|\text{Aut}(K)| > 2$. Actually $\text{Aut}(K) = V_4$. We verified this by using equation (17) from [13] which gives a necessary and sufficient condition for a genus 2 field K to have $V_4 \leq \text{Aut}(K)$ (in terms of the classical invariants of K). Taking the resultant of this equation and the one in Appendix A yields an expression that has (22) (the projectivized version) as a factor. From the conditions given in [13] for K to have $\text{Aut}(K)$ strictly bigger than V_4 , one checks that actually $\text{Aut}(K) = V_4$ for all K in locus (22).*

4.1. The general case. By previous section θ is generically a covering of degree 2. So there exists a Zariski open subset $\mathcal{U}$ of k^2 with the following properties: Firstly, θ is defined everywhere on $\mathcal{U}$ and is a covering of degree 2 from $\mathcal{U}$ to $\theta(\mathcal{U})$. Further, if $\mathbf{u} \in \mathcal{U}$ then all $\mathbf{u}' \in k^2$ with θ defined at $\mathbf{u}'$ and $\theta(\mathbf{u}') = \theta(\mathbf{u})$ also lie in $\mathcal{U}$. Suppose $\underline{i} \in k^3$ such that $|\theta^{-1}(\underline{i})| > 2$ and $\det(\text{Jac}(\theta))$ does not vanish at any point of $\theta^{-1}(\underline{i})$. Then by implicit function theorem, there is an open ball B around each element of $\theta^{-1}(\underline{i})$ such that each point in $\theta(B)$ has > 2 inverse images under θ . But B has to intersect the Zariski open set $\mathcal{U}$. This is a contradiction. Thus, if $\underline{i} \in k^3$ and $|\theta^{-1}(\underline{i})| > 2$, then $\det(\text{Jac}(\theta)) = 0$ at some point of $\theta^{-1}(\underline{i})$ and so $\underline{i}$ satisfies (22). It follows that if K is a genus 2 field with $J_2 \neq 0$ and $G \neq 0$ then $e_3(K) = 2$.

Note that if there is a degenerate pair (K, E_1) and a non-degenerate pair (K, E_2) (for the same K) then for the (u, v) -invariants of the latter we have (12), hence $\beta(v) = 27$ and so the β -image of (u, v) doesn't correspond to a non-degenerate pair.

We can now complete the proof of part (ii) of Main Theorem. We noted before that the condition in (ii) is necessary. This condition (see Appendix A) defines an irreducible sublocus of $\mathcal{M}_2$ because the corresponding equation is irreducible (by Maple). The locus $e_3(K) \geq 1$ lies in this sublocus and contain a dense subset of it (image of θ). By Lange [3], the locus $e_3(K) \geq 1$ is Zariski closed, hence equals the locus defined by the equation in Appendix A.

4.2. Exceptional Cases for $J_2 = 0$. We consider the case when $J_2 = 0$, where

$$(23) \quad \begin{aligned} J_2 &= -2(3v^2 + 4u^2 - 12uv + 252u - 54v - 405) \\ &= \frac{2}{3}(3v - 6u + 2u\sqrt{6} - 27 - 18\sqrt{6})(3v - 27 + 18\sqrt{6} - 6u - 2u\sqrt{6}) \end{aligned}$$

Genus 2 fields K with $J_2 = J_4 = 0$ and $J_6 \neq 0$ (resp., $J_2 = J_6 = 0$ and $J_4 \neq 0$) are classified up to isomorphism by the invariant $\frac{J_5^5}{J_{10}^5}$ (resp., $\frac{J_5^5}{J_{10}^5}$), see Igusa [9]. In the first (resp., second) case there are exactly two (resp., four) such K with $e_3(K) \geq 1$ and they all have $e_3(K) = 2$.

The invariants

$$(24) \quad a_1 := \frac{J_4 \cdot J_6}{J_{10}}, \quad a_2 := \frac{J_6 \cdot J_{10}}{J_4^4}$$

determine genus two fields with $J_2 = 0$, $J_4 \neq 0$, and $J_6 \neq 0$ up to isomorphism, see Igusa [9]. A field K with such invariants has $e_3(K) \geq 1$ if and only if

$$(25) \quad \begin{aligned} & 46656a_1^5a_2^3 + 7558272a_1^4a_2^3 + 1259712\sqrt{6}a_1^4a_2^3 - 15552a_1^4a_2^2 - 12427478784a_1^3a_2^2 + 4917635712\sqrt{6}a_1^3a_2^2 \\ & + 1728a_1^3a_2 - 656217531654480a_1^2a_2^2 + 267571209034080\sqrt{6}a_1^2a_2^2 - 1844125056a_1^2a_2 + 743525568\sqrt{6}a_1^2a_2 \\ & - 64a_1^2 - 6334497449472117312a_1a_2^2 + 2585860435265558832\sqrt{6}a_1a_2^2 + 230833239838992a_2a_1 - \\ & 94237227087840a_2a_1\sqrt{6} - 601244429975805030777a_2^2 + 245429539257764380572a_2^2\sqrt{6} = 0 \end{aligned}$$

or

$$(26) \quad \begin{aligned} & 230833239838992a_2a_1 + 94237227087840a_2a_1\sqrt{6} - 12427478784a_1^3a_2^2 - 4917635712\sqrt{6}a_1^3a_2^2 \\ & + 1728a_1^3a_2 - 15552a_1^4a_2^2 - 656217531654480a_1^2a_2^2 - 267571209034080\sqrt{6}a_1^2a_2^2 + 46656a_1^5a_2^3 - 64a_1^2 \\ & - 601244429975805030777a_2^2 - 245429539257764380572a_2^2\sqrt{6} - 1844125056a_1^2a_2 - 743525568\sqrt{6}a_1^2a_2 \\ & - 6334497449472117312a_1a_2^2 - 2585860435265558832\sqrt{6}a_1a_2^2 + 7558272a_1^4a_2^3 - 1259712\sqrt{6}a_1^4a_2^3 = 0 \end{aligned}$$

As one can check by discussing the map

$$\vartheta : (u, v) \rightarrow (a_1, a_2)$$

these fields K generically have $e_3(K) = 2$. It remains to check the singular points of the curves (25) and (26). The intersection of the two components is empty. Each component has four singular points two of which have $e_3(K) = 2$ and the other two have $e_3(K) = 4$. Thus there are exactly four genus 2 fields with $e_3(K) = 4$. Two of them satisfy (25), we call them P_1 and P_2 . The other two are obtained from P_1, P_2 by applying the automorphism $\sqrt{6} \rightarrow -\sqrt{6}$.

P_1	$a_1 = -\frac{77169}{8} + \frac{30759}{8}\sqrt{6}, \quad a_2 = \frac{13783592}{23149125} + \frac{5629912}{23149125}\sqrt{6}$
P_2	$a_1 = -\frac{650835}{8} + \frac{268785}{8}\sqrt{6}, \quad a_2 = -\frac{2984}{20002028625} - \frac{144872}{60006085875}\sqrt{6}$

TABLE 2. Singular points of (25) with $e_3(K) = 4$

The four values of (u, v) corresponding to P_1 (resp., P_2) are obtained as follows: u is determined by v via

$$u = \frac{1}{2}(3 + \sqrt{6})(v - 9 - 6\sqrt{6})$$

and v is one of the 4 roots of

$$(27) \quad \begin{aligned} & 9v^4 + (-693 - 27\sqrt{6})v^3 + (15141 + 1749\sqrt{6})v^2 + (-66414 - 33396\sqrt{6})v \\ & - 1368\sqrt{6} + 3388 = 0 \end{aligned}$$

(resp.)

$$(28) \quad \begin{aligned} & 3v^4 + (-153 + 33\sqrt{6})v^3 + (3129 - 1219\sqrt{6})v^2 + (-10854 - 11556\sqrt{6})v \\ & + 176904\sqrt{6} + 450036 = 0 \end{aligned}$$

This completes the proof of the Main Theorem. $\square$

5. J-INVARIANTS

A genus 2 field K corresponding to a generic point of locus $e_3(K) \geq 1$ has exactly 2 elliptic subfields E_1 and E_2 of degree 3. We can take $E_1 = E$ from lemma 3. Its j -invariant is

$$(29) \quad j_1 = 16v \frac{(vu^2 + 216u^2 - 126vu - 972u + 12v^2 + 405v)^3}{(v-27)^3(4v^2 + 27v + 4u^3 - 18vu - vu^2)^2}$$

The automorphism $\beta \in \text{Gal}_{k(u,v)/k(r_1, r_2)}$ permutes E_1 and E_2 , therefore switches j_1 and j_2 . Then $j_2 = \beta(j_1)$ is

$$(30) \quad j_2 = -256 \frac{(u^2 - 3v)^3}{v(4v^2 + 27v + 4u^3 - 18vu - vu^2)}$$

For the equation of E_2 using a different approach see [12], chapter 5.

5.1. Isomorphic Elliptic Subfields. Suppose that $E_1 \cong E_2$. Then, $j_1 = j_2$ implies that

$$(31) \quad 8v^3 + 27v^2 - 54uv^2 - u^2v^2 + 108u^2v + 4u^3v - 108u^3 = 0$$

or

$$(32) \quad \begin{aligned} & 324v^4u^2 - 5832v^4u + 37908v^4 - 314928v^3u - 81v^3u^4 + 255879v^3 + 30618v^3u^2 \\ & - 864v^3u^3 - 6377292uv^2 + 8503056v^2 - 324u^5v^2 + 2125764u^2v^2 - 215784u^3v^2 \\ & + 14580u^4v^2 + 16u^6v^2 + 78732u^3v + 8748u^5v - 864u^6v - 157464u^4v + 11664u^6 = 0 \end{aligned}$$

Remark 11. *The former equation is the condition that $\det(\text{Jac}(\theta)) = 0$ see equation (21).*

Remark 12. *If $e_3(K) \geq 1$ then the automorphism group $\text{Aut}(K/k)$ is one of the following: $\mathbb{Z}_2, V_4, D_8$, or D_{12} . $\mathbb{Z}_2$ is the generic case, V_4 is a 1-dimensional family. There are exactly 6 genus 2 fields with automorphism group D_8 (resp., D_{12}).*

For the proof see [12].

Appendix A:

Here is the equation which defines the locus $\mathcal{L}_3$ of fields K with $e_3(K) \geq 1$.

$$(33) \quad C_8 J_{10}^8 + C_7 J_{10}^7 + \cdots + C_1 J_{10} + C_0 = 0$$

where $C_0, \dots, C_8$ are

$$\begin{aligned} C_8 &= 2^4 \cdot 3^{31} \cdot 5^5 \cdot 19^{10} \cdot 29^5 \\ C_7 &= 2^4 \cdot 3^{27} \cdot 19^5 (-194894640029511 J_2^5 - 55588661819356000 J_4^2 J_2 - 12239149540657725 J_2^3 J_4 \\ &+ 223103526505680000 J_4 J_6 + 40811702108053500 J_2^2 J_6) \\ C_6 &= 2^2 \cdot 3^{21} (-35802284468757765858432 J_4^5 - 1756270399106587730391 J_4^2 J_2^6 - 28638991859523006654 J_4 J_2^8 \\ &- 84091225203760159441286 J_4^3 J_2^4 + 400895959391006953561032 J_4^4 J_2^2 - 61773685738999443 J_2^1 0 \\ &- 3673201396072259603756160 J_4^3 J_2 J_6 + 7879491755218264984387200 J_4^2 J_6^2 + 15251447355608658629952 J_2^5 J_4 J_6 \\ &+ 1179903008384844066250272 J_2^3 J_4^2 J_6 - 5566672398589809889658760 J_2^2 J_4 J_6^2 + 112024289372554183680 J_2^7 J_6 \\ &- 32116769409722716182888 J_2^4 J_6^2 + 851217187754962249155200 J_2 J_6^3) \end{aligned}$$

$$\begin{aligned}
C_5 = & 2^2 \cdot 3^{19} (-12630004382695462653 J_4^4 J_2^7 + 320839252764287362560 J_4^7 J_2 - 1876069272397136886448 J_4^6 J_2^3 \\
& + 606742866220456356580 J_4^5 J_2^5 - 124173485719052715 J_4^3 J_2^9 + 22241034512101438944000 J_4^5 J_2^2 J_6 \\
& - 88546736703938826304512 J_4^4 J_2 J_6^2 - 10712078926420753449984 J_2^4 J_4 J_6 + 68904635323303664511264 J_2^3 J_4^3 J_6^2 \\
& - 192353895694677164016384 J_2^2 J_4^2 J_6^3 + 197449733923926905783808 J_2 J_4 J_6^4 - 1916173047371645223936 J_4^6 J_6 \\
& + 116211018774997425051648 J_4^3 J_6^3 - 132143597697786172416 J_6^5 + 1361403542457288 J_2^1 J_4 J_6 \\
& - 5005765118740492656 J_2^2 J_4 J_6^2 + 232819061639483430720 J_2^6 J_4^3 J_6 - 1576319894694585178452 J_2^5 J_4^2 J_6^2 \\
& + 4655239459208764553088 J_2^4 J_4 J_6^3 - 226900590409548 J_2^{11} J_4^2 - 2042105313685932 J_2^9 J_6^2 + 6261632755967872800 J_2^6 J_6^3 \\
& - 5065734796478275576176 J_2^3 J_6^4 + 1352318109350828796 J_2^8 J_4^2 J_6)
\end{aligned}$$

$$\begin{aligned}
C_4 = & 3^{15} (1417825317153277312 J_4^9 J_2^2 + 2391308818408811717 J_4^6 J_2^8 + 718590303030600 J_2^{10} J_4^5 \\
& - 638760745337170544640 J_4 J_6^6 + 440759275303802880 J_4^{10} - 8118717280771686540192 J_4^5 J_4^2 J_6^2 \\
& + 42668434906863398019072 J_2^4 J_2 J_6^5 - 57054664814020640574336 J_4^3 J_2^2 J_6^4 + 30546774740158581676032 J_4^4 J_2^3 J_6^3 \\
& + 8601814215123831275904 J_4^6 J_2^2 J_6^2 - 1449562700682195916800 J_4^7 J_2^3 J_6 + 1067928354124249303104 J_4^8 J_2^5 J_6 \\
& - 10443896263024316301312 J_2^3 J_4 J_6^5 + 7247970315150439028112 J_2^4 J_4^4 J_6 - 21769241176751736619008 J_2^5 J_4^2 J_6^3 \\
& - 2640201919999154595648 J_2^5 J_4^3 J_6^3 - 55893562424445261312 J_2^7 J_4^5 J_6 + 531409635241191119304 J_2^6 J_4^4 J_6^2 \\
& - 1012614205133520 J_2^8 J_4^4 J_6 - 2454855015326199552 J_2^5 J_6^5 - 12501409939920 J_2^{12} J_4^4 - 675076136755680 J_2^{10} J_4^2 J_6^2 \\
& - 33390518666828400 J_2^9 J_4^4 J_6 + 3188363568027498432 J_2^6 J_4 J_6^4 - 1569001498547402304 J_2^7 J_4^2 J_6^3 \\
& - 275375222428239820800 J_2^7 J_6^2 + 19809849095518050330624 J_4^4 J_6^4 + 6179516061983740183680 J_2^2 J_6^6 \\
& + 150016919279040 J_2^{11} J_4^3 J_6 + 87799481406335621136 J_4^8 J_2^4 + 1350152273511360 J_2^9 J_4 J_6^3 \\
& - 55496611186132800648 J_4^7 J_6^6 + 39911809855842557952 J_4^8 J_2 J_6 + 353362680242481096 J_2^8 J_4^3 J_6^2)
\end{aligned}$$

$$\begin{aligned}
C_3 = & 2^4 \cdot 3^{12} (-19225816442103600 J_4^{10} J_2^5 + 6433952690394144 J_2^4 J_6^7 - 2917203075615 J_2^{11} J_4^7 \\
& + 62951605613640 J_2^{10} J_4^6 J_6 + 7900854051362368 J_4^{11} J_2^3 - 873165547551982 J_2^9 J_4^8 + 13234982161044480 J_4^{12} J_2 \\
& + 4077902864550187008 J_4^5 J_6^5 + 7506792545698293 J_4^9 J_2^7 - 55019014994202624 J_4^{11} J_6 - 3415519987075510272 J_4^7 J_6^7 \\
& - 932605137272623104 J_4^8 J_6^3 - 6607177263254292480 J_2 J_6^8 - 1394785406520 J_2^7 J_6^6 - 1913285880 J_2^{13} J_4^6 \\
& - 258293593800 J_2^{11} J_4^4 J_6^2 - 1976299597616301504 J_4^8 J_2^3 J_6^2 + 1337598192058041744 J_4^7 J_2^5 J_6^2 \\
& - 2324642344200 J_2^9 J_4^4 J_6 + 2789570813040 J_2^8 J_4 J_6^5 - 243015467955111198 J_2^7 J_4^6 J_6^2 + 22136761801348668 J_2^8 J_4^7 J_6 \\
& - 155463896437263612 J_4^8 J_2 J_6 + 16101033796183004352 J_4^5 J_2^3 J_6^4 + 16367298631796450304 J_4^3 J_2 J_6^6 + 8433152 J_4^7 J_2^2 J_6^3 \\
& - 8254965178021469184 J_6^4 J_2 J_6^4 + 34439145840 J_2^{12} J_4^5 J_6 - 576988130682378 J_2^9 J_4^5 J_6^2 + 2912934238489260 J_2^8 J_4^4 J_6^3 \\
& - 8749875412454175 J_2^7 J_4^4 J_6^4 + 15637511592200340 J_2^6 J_4^2 J_6^5 - 127105068829245696 J_4^{10} J_2 J_6 + 614908581517421568 J_4^9 J_2 J_2^2 \\
& - 23374419431360207616 J_4^4 J_2^2 J_6^5 + 1508868948605946984 J_2^6 J_4^5 J_6^3 - 5795040294470623824 J_2^5 J_4^4 J_6^4 \\
& + 14094983896511630112 J_2^4 J_4^3 J_6^5 + 1033174375200 J_2^{10} J_4^3 J_6^3 + 314069798204069472 J_2^9 J_4^4 J_6 - 61501104 J_4^6 J_2^4 J_6^3 \\
& - 21194163800222025024 J_2^3 J_4^2 J_6^6 + 18002402119176332544 J_2^2 J_4 J_6^7 - 15392091937240080 J_2^5 J_4 J_6^6)
\end{aligned}$$

$$\begin{aligned}
C_2 = & 2^5 \cdot 3^8 (-159732958548480 J_4^{13} J_2 J_6 - 27945192968593920 J_2 J_4 J_6^9 + 238596124150086 J_4^{10} J_2^7 J_6 + 3224288 J_2^3 J_6^9 \\
& - 36311136215244 J_2^9 J_4^9 J_6 - 996173640 J_2^{12} J_4^6 J_6^2 + 5977041840 J_2^{11} J_4^5 J_6^3 - 22413906900 J_2^{10} J_4^4 J_6^4 - 37501414009508 J_4^{13} J_2^4 \\
& + 14210312049697149 J_2^6 J_4^6 J_6^4 + 86354918885580768 J_2^4 J_4^6 J_6^6 - 111444977082978432 J_2^3 J_4^3 J_6^7 + 27908893977856 J_4^{14} J_2^2 \\
& + 83768141083825152 J_2^2 J_4^2 J_6^8 - 42942980968765488 J_2^5 J_4^5 J_6^5 - 10736445647473 J_4^{11} J_2^8 + 61746352553318400 J_4^4 J_2 J_6^7 \\
& + 410958880454688 J_4^{12} J_2^3 J_6 - 14059252057660416 J_4^3 J_6^8 - 1643659809866496 J_4^{11} J_2^2 J_6^2 + 441832778741790 J_2^8 J_4^8 J_6^2 \\
& - 3128599551108636 J_2^7 J_4^7 J_6^3 + 2815950495430656 J_4^{10} J_2 J_6^3 + 12291244885171152 J_4^8 J_2^5 J_6^3 + 1579225145 J_2^{12} J_4^9 \\
& - 2286353789913249 J_4^9 J_2^6 J_6^2 - 40300476525629352 J_4^7 J_2^4 J_6^4 + 81707043798929088 J_4^6 J_2^3 J_6^5 - 25936092270 J_2^6 J_6^8 \\
& - 98257765274489088 J_4^5 J_2^2 J_6^6 + 3318887207480832 J_4^{10} J_2^4 J_6^2 - 478511899451856 J_4^{11} J_2^5 J_6 + 28476287051677 J_4^{12} J_2 J_6^2 \\
& - 12153253649302656 J_4^9 J_2^3 J_6^3 + 24757975700165376 J_4^8 J_2^2 J_6^4 - 26570902457981952 J_4^7 J_2 J_6^5 - 6755065089024 J_4^{15} \\
& + 10883911680 J_4^{10} + 53793376560 J_2^9 J_4^3 J_6^5 - 80690064840 J_2^8 J_4^2 J_6^6 + 69162912720 J_2^7 J_4 J_6^7 + 94873680 J_2^{13} J_4^7 J_6 \\
& + 227109129291 J_2^{10} J_4^7 J_6^2 - 628213747356 J_2^9 J_4^6 J_6^3 - 1389130574661 J_2^8 J_4^5 J_6^4 + 16465793988870 J_2^7 J_4^4 J_6^5 \\
& - 56794191944715 J_2^6 J_4^3 J_6^6 + 102713329135152 J_2^5 J_4^2 J_6^7 - 98529746457492 J_2^4 J_4 J_6^8 - 30650938650 J_2^{11} J_4^8 J_6 \\
& - 1716480768 J_2^9 J_4^6 J_6 + 11718053954519040 J_2^6 J_4^6 J_6^6 + 220752428322816 J_2^{12} J_6^2 + 1322792799725 J_2^{10} J_4^{10} - 3953070 J_2^{14} J_4^8)
\end{aligned}$$

$$\begin{aligned}
C_1 = & -2^8 \cdot 3^5 (61736960 J_4^8 J_2 - 182135808 J_4^7 J_6 + 16021872 J_4^7 J_2^3 - 211022400 J_4^6 J_2^2 J_6 - 26594919 J_4^6 J_2^5 \\
& + 899159040 J_4^5 J_2^2 J_6^2 + 330458928 J_4^5 J_2 J_6 - 215198 J_2^7 J_4^5 - 1227405312 J_4^4 J_6^3 - 1535734368 J_4^4 J_2^3 J_6^2 \\
& + 2930532 J_4^4 J_2^2 J_6 - 363 J_2^9 J_4^4 + 3162070656 J_4^3 J_2^2 J_6^3 - 16471998 J_4^3 J_2^5 J_6^2 + 4356 J_4^3 J_2^8 J_6 - 19602 J_4^2 J_2^7 J_6^2)
\end{aligned}$$

$$- 2433162240J_4^2J_2J_6^4 + 47961936J_4^2J_2^4J_6^3 + 39204J_4J_2^6J_6^3 - 72369936J_4J_2^3J_6^4 + 746496J_4J_6^5 - 29403J_2^5J_6^4 \\ + 45116352J_2^2J_6^5)(J_4^3 - J_2^2J_4^2 + 6J_2J_6J_4 - 9J_6^2)^3$$

$$C_0 = 2^8(768J_4^2 - 416J_4J_2^2 - J_2^4 + 1536J_2J_6)(J_4^3 - J_2^2J_4^2 + 6J_2J_6J_4 - 9J_6^2)^6$$

Appendix B:

The equation of the branch locus of the map

$$\theta : k^2 \setminus \{\Delta = 0\} \rightarrow \mathcal{L}_3 \\ (u, v) \rightarrow (i_1, i_2, i_3)$$

$$(34) \quad \begin{aligned} & 3507505273398025i_1^{12} - 4880484817793862073548480i_1^{11} + 192(11302504938388489628125i_2 \\ & + 346452039237689650581192)i_1^{10} - 4(89439905046278964319358976 + 60307157046030532997225i_2^2 \\ & - 4188981066069113234648640i_2)i_1^9 + 192(4924672355809568004885504 + 33588887753890413143515i_2^2 \\ & - 89290373874540245356608i_2)i_1^8 - 192(6360139591235383381327872 + 593587805135845078438632i_2^2 \\ & + 849397678885114696829952i_2 - 15537701670163340329775i_2^3)i_1^7 + 2(406606074742962841916289024i_2^2 \\ & - 27711519511099200420652160i_2^3 + 305526006347596356290347008 + 183049808606955774794075i_2^4 \\ & + 255320765313220782576893952i_2)i_1^6 - 192i_2(13654023711946692280289280i_2 + 60377228453350507376315i_2^3 \\ & - 1464350022771442997265792i_2^2 + 2245754530466537929703424)i_1^5 + 192i_2^2(19906166147692916436664320 \\ & + 524465846991778455117432i_2^2 - 2333360138590692031481856i_2 - 10694743521252049023175i_2^3)i_1^4 \\ & + 20i_2^2(2535928320048654288481984i_2^3 - 5763664926963183845376000i_2 - 28262691566657845506883584i_2^2 \\ & - 99413657833844087193600000 - 9280304409903257402325i_2^4)i_1^3 - 192i_2^3(1807602326421361731982656i_2^2 \\ & - 76401752022567738591625i_2^3 - 2867627019656613888000000 - 9918952715424066995911680i_2)i_1^2 \\ & + 1728i_2^4(1262371230708245434125i_2^3 - 44970318919363276752280i_2^2 - 1752882366993587712000000 \\ & + 506870449554602235884544i_2)i_1 + i_2^4(127385395640432909375625i_2^4 - 6059765135286837968377600i_2^3 \\ & - 717148324858373259264000000i_2 + 102838387547772598641352704i_2^2 + 1617386925920256 \cdot 10^{12}) = 0 \end{aligned}$$

REFERENCES

- [1] G. FREY, On elliptic curves with isomorphic torsion structures and corresponding curves of genus 2. *Elliptic curves, modular forms, and Fermat's last theorem (Hong Kong, 1993)*, 79-98, Ser. Number Theory, I, *Internat. Press, Cambridge, MA*, (1995).
- [2] G. FREY AND E. KANI, Curves of genus 2 covering elliptic curves and an arithmetic application. *Arithmetic algebraic geometry (Texel, 1989)*, 153-176, *Progr. Math.*, 89, *Birkhäuser Boston, Boston, MA*, (1991).
- [3] H. LANGE, Über die Modulvarietät der Kurven vom Geschlecht 2. *J. Reine Angew. Math.*, 281, 80-96, 1976.
- [4] E. KANI AND W. SCHANZ, Diagonal quotient surfaces. *Manuscripta Math.* 93 (1997), no. 1, 67-108.
- [5] E. KANI, The number of curves of genus two with elliptic differentials. *J. Reine Angew. Math.* 485 (1997), 93-121.
- [6] A. KRAZER, *Lehrbuch der Thetafunctionen*, Chelsea, New York, (1970).
- [7] V. KRISHNAMOORTHY, Invariants of genus 2 curves, *PhD thesis*, University of Florida, 2001.
- [8] M. R. KUHN, Curves of genus 2 with split Jacobian. *Trans. Amer. Math. Soc* **307** (1988), 41-49
- [9] J. IGUSA, Arithmetic Variety Moduli for genus 2. *Ann. of Math.* (2), 72, 612-649, (1960).
- [10] MAPLE 6, Waterloo Maple Inc., 2000.
- [11] T. SHASKA, Genus 2 Curves With (N,N) Decomposable Jacobians, *Jour. Symb. Comp.*, Vol 31, **no. 5**, pg. 603-617, 2001.

- [12] T. SHASKA, Curves of Genus Two Covering Elliptic Curves, *PhD thesis*, University of Florida, 2001.
- [13] T. SHASKA AND H. VÖLKLEIN, Elliptic Subfields and automorphisms of genus 2 function fields, to appear in: *Proceeding of the Conference on Algebra and Algebraic Geometry with Applications: The celebration of the seventieth birthday of Professor S.S. Abhyankar*, Springer-Verlag, 2000.
- [14] G. TAMME, Ein Satz über hyperelliptische Funktionenkörper. J. Reine Angew. Math. 257 (1972), 217–220.
- [15] H. VÖLKLEIN, Groups as Galois Groups – an Introduction, Cambr. Studies in Adv. Math. 53, Cambridge Univ. Press 1996.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT IRVINE, IRVINE, CA 92697.
E-mail address: `tshaska@math.uci.edu`